Precipitation.—The average was 0.37 below the normal; greatest monthly, 2.25, at Oelrichs; least monthly, trace, at a number of stations.

Wind.—Prevailing direction, northwest.—S. W. Glenn, Local Forecast Official, Weather Bureau, Huron, director.

TENNESSEE WEATHER AND CROP SERVICE.

Temperature.—The mean was 3.7 below the normal; maximum, 73, at Chattanooga, 9th; minimum, 4, at Trenton, 26th; greatest monthly range, 61, at Trenton; least monthly range, 43, at Springdale.

Precipitation.—The average was 3.80 above the normal; greatest monthly,

12.14, at Covington; least monthly, 3.70, at Strawberry Plains.

Wind.—Prevailing direction, north.—J. B. Marbury, Local Forecast Official, Weather Bureau, Nashville. director.

TEXAS.

Fort Ringgold, 12th; minimum, -9, at Coldwater, 12th; greatest monthly range, 77, at Fort Hancock; least monthly range, 43, at McGregor.

recipitation.—The average was 0.76 below the normal; greatest monthly, 6.61, at Orange; least monthly, 0.00, at Forts Clark and Hancock.

Wind.—Prevailing direction, north.—D. D. Bryan, Galveston, director; I. M. Cline, Local Forecast Official, Weather Bureau, assistant.

Temperature. - Maximum, 68, at Saint George, 27th; minimum, -34, at Randolph, 23d; greatest monthly range, 75, at Thistle; least monthly range, 37, at Salt Lake City.

Precipitation.—Greatest monthly, 4.70, at Silver Lake; least monthly, 0.19,

at Fort Du Chesne.

Wind. — Prevailing direction, northwest. — G. N. Salisbury, Observer, Weather Bureau, Salt Lake City, director.

(See New England.)

VIRGINIA.

Temperature.—Maximum, 74, at Norfolk, 10th, and at Richmond, 9th; minimum, 7, at Hot Springs and Warsaw, 25th; greatest monthly range, 62, at Warsaw; least monthly range, 47, at Cape Charles.

Precipitation.—Greatest monthly, 7.50, at Birdsnest; least monthly, 2.25, at Christiansburg

Wind.—Prevailing direction, north.—Dr. E. A. Craighill, Lynchburg, director: J. N. Ryker. Observer. Weather Bureau. assistant.

WASHINGTON

Temperature.—The mean was 3.9 below the normal; maximum, 61, at Fort Simcoe, 28th; minimum, -17, at Pullman, 22d; greatest monthly range, 65, at Ellensburg; least monthly range, 23, at Tatoosh Island and Fort Canby. Precipitation.—The average was 0.57 below the normal; greatest monthly, 16.52, at Neah Bay; least monthly, 0.32, at Davenport.

Wind.—Prevailing direction, south.—H. F. Alciatore, Observer, Weather

Bureau, Seattle, director.

WEST VIRGINIA.

Temperature.—Maximum, 74, at Point Pleasant, 9th; minimum, -5, at Temperature.—The mean was 2.2 below the normal; maximum, 94, at | Pleasant Hill, 24th; greatest monthly range, 65, at Pleasant Hill; least monthly range, 48, at Elkhorn.

Frecipitation.—Greatest monthly, 7.48, at Beverly; least monthly, 2.05, at Cloverdale.

Wind .- Prevailing direction, west.-H. W. Richardson, Observer, Weather Bureau, Parkersburg, director.

WISCONSIN.

Temperature.—The mean was 1.5 below the normal; maximum, 60, at Whitehall, 28th; minimum, —35, at Butternut, 24th; greatest monthly range, 83, at Butternut; least monthly range, 48, at Madison.

Precipitation.—The average was 0.50 below the normal; greatest monthly. 1.95, at Sharon; least monthly, 0.15, at Osceola.

Wind.—Prevailing directions, southwest and northwest.—W. L. Moore, Local Forecast Official, Weather Bureau, Milwaukee, director.

Temperature. - Maximum, 64, at Wheatland, 27th; minimum, -32, at Sheridan, 21st; greatest monthly range, 78, at Sheridan; least monthly range, 65, at Fort Yellowstone.

Precipitation.—The average was 0.09 above the normal; greatest monthly,

2.30, at Labarge; least monthly, 0.10, at Laramie.

Wind.—Prevailing direction, west.—E. M. Ravenscraft. Observer, Weather Bureau, Cheyenne, director.

INLAND NAVIGATION.

STAGE OF WATER IN RIVERS.

The following table shows the danger point at each river station; the highest and lowest stages for the month of February, 1894, with the dates of occurrence, and the monthly range:

Heights of rivers above low-water mark, February, 1894.

Stations.	Danger- point on gauge.	Highest water.		Lowest water.		outhly range.
		Height.	Date.	Height.	Date.	Mon
Red River.	Feet.	Feet.		Feet.		Feet.
Bhreveport, La	29.2	18.9	28	5-4	6	13.5
Fort Smith, Ark	22.0	13.8	13	3.2	I, 2	10.6
Little Rock, Ark	23.0	21.0	13	6.5	2, 3	14.5
Pierre, S. Dak *	13.0					
Bioux City, Iowa *	18.7				· · · · · · · · · · · · · · · · · · ·	
Kansas City, Mo†	21.0	8.8	25	l i	17	2.2
t. Paul, Minn *	14-0				• • • • • • • • • • • •	• • • • • •
La Crosse, Wis *	10.0					
Dubuque, Iowa *	16.0					
Davenport, Iowa *	15-0					• • • • • •
Keokuk, Iowa*	14.0			· · · · · · ː · ·		*****
Hannibal, Mo	17.0	3.4		0.8	13	2.6
St. Louis, Mo	30.0	6.4	II	0.2	3	6.3
Cairo, Ill	40.0	37.0		13.4	4	23.6
Memphis, Tenn	33.0	29.4	20	9.6	. 4	19.8
Vicksburg, Miss	41.0	40.3	27	16.3	7, <u>8</u>	24.0
New Orleans, La	13.0	13.2	27, 28	5.7		7.5
Parkersburg, W. Va	38.0	24.0	13	8-6	4	15.4
Cincinnati, Ohio	45.0	35.6	15	17.2	4	18.4
Louisville, Ky	24.0	12.9	14	7-7	I	5.2
Nashville, Tenn	40.0	41.9	10	9.3	1	32.6
Chattanooga, Tenn	33.0	25.5	6	4-9	3	20.6
Knoxville, Tenn	29.0	15.4	5	2.3	2	13.1
Pittsburg, Pa	22.0	14.8	11	3.8	27	11.0
Augusta, Ga	32.6	20.9	16	7.8	4, 10	13.1
Portland, Oregon	15.0	8-4	11	4.2	27	4.2

Heighte of viners Continued

Stations.	Danger- point on gauge.	Highest water.		Lowest water.		thly ge.
		Height.	Date.	Height.	Date,	Monthly range.
Susquehanna River.	Feet.	Feet.		Feet.		Feet.
Harrisburg, Pa	17.0	6.0	12	2.0	6-8	4.0
Montgomery, Ala	48.0	27 - 1	16	3.5	3	23.6
Lynchburg, Va	18.0	5.3	- 5	1.5	Ι, 2	3.8
Red Bluff, Cal	22.0	13.0	20	4.0	14, 17	9.0
Sacramento, Cal Des Moines River.	25.0	22.5	21, 22	18.5	16	4.0
Des Moines, Iowa*	19.0		• • • • • • • • • • • • • • • • • • • •	-		
*Frozen.		<u> </u>	† For 15	days.		·

† For 15 days.

FLOODS.

The above table shows that in the rivers here recorded the nearest approaches to the danger line have been at Little Rock, Ark., on the 13th; Cairo, Ill., on the 16th; Sacramento, Cal., on the 21st and 22d; New Orleans, La., on the 27th and 28th.

The only notable flood was in the Cumberland River which reached the danger line on the morning of the 9th, attained its maximum on the 10th, and was below the danger line on the morning of the 11th.

NAVIGATION OF RIVERS AND HARBORS.

The weekly "bulletin of snow lying on the ground" gives reports of ice in rivers and harbors, and shows that on Monday, February 5, the ice in the Mississippi River varied from 10 inches at Hannibal, Mo., to 20 or 22 inches at La Crosse, Wis., and St. Paul, Minn.; in the Missouri River it varied from 3 inches at Kansas City, Mo., to 30 inches at Bismarck, N.

Dak., and 24 inches at Williston, N. Dak. On the Great Lakes there were from 4 to 6 inches in the harbors of lakes Ontario, Erie, and Michigan, 8 inches on Lake Huron, and 22 inches on Lake Superior, at Duluth, Minn. These conditions remained with but little change throughout the month. On Monday, the 26th, the reported thickness of ice, in inches, was as follows:

Hudson River.—Albany, N. Y., 14.

Upper Mississippi River.—St. Paul, Minn., 24; La Crosse, Wis., 28; Dubuque, Iowa, 12; Davenport, Iowa, 12; Keokuk, Iowa, 12; Hannibal, Mo., 6.

Missouri River.—Williston, N. Dak., 42; Bismarck, N. Dak., 28; Pierre, S. Dak., 29; Sioux City, Iowa, 18; Omaha, Nebr.,

14; Kansas City, Mo., 4.

Arkansas River.—Fort Smith, Ark., 10; Dodge City, Kans., 2. Platte River.—North Platte, Nebr., 16.5. Red River of the North.—St. Vincent, Minn., 24.

Atlantic Coast.—Block Island, R. I., 3; New Haven, Conn.,

4; Portland, Me., 3.

Lake Ontario.—Oswego, N. Y., 9.5; Rochester, N. Y., 6.5. Lake Erie.—Buffalo, N. Y., 10; Erie, Pa., 10; Cleveland, Ohio, 6; Sandusky, Ohio, 8; Toledo, Ohio, 6.3; Detroit, Mich., 9.5.

Lake Huron.—Port Huron, Mich., 9.5; Alpena, Mich., 16. Lake Michigan.—Grand Haven, Mich., 11; Chicago, Ill., 4; Green Bay, Wis., 15.

Lake Superior.—Sault Ste. Marie, Mich., 36; Marquette, Mich., 9; Duluth, Minn., 29.5.

The following special reports have also been collected:

Lake Superior. - Marquette, Mich., 21st, ice moving out of harbor; 22d, harbor clear of ice, except back of breakwater.

Lake Michigan. - Grand Haven, Mich., 20th, for the first time during the winter ice is obstructing navigation; slush ice fills the harbor and extends about 5 miles out into the lake. The steamer Wisconsin remained in port, unable to force her way through; 21st, the steamer Roanoke is reported fast in the ice 3 miles out; 24th, southeast winds cleared the harbor of ice and enabled the steamer Roanoke to enter port; 27th, immense floes of ice cover the lake, and the Milwaukee boat failed to make her way through and returned to port; 28th, navigation stopped by immense fields of ice.

Lake Huron. -Port Huron, Mich., 1st to 5th, floating ice in the St. Clair River; 6th, an ice gorge formed in the St. Clair at Marysville, 6 miles south of Port Huron; 9th and 11th, floating ice in the St. Clair; 22d, an ice gorge

ormed one half mile below the city.

Missouri River.—Hermann, Mo., gorge broken on the 7th and ice moving out; 12th, heavy floating ice; 15th, frozen; 18th, opened again; 21st, heavy floating ice.

Hilmois River.—Ottawa, Ill., 9th, ice passed out.

Ohio River.—Marietta, Ohio, 1st to 9th, heavy floating ice; 17th to 18th, floating ice.* Parkersburg, W. Va., 1st to 6th, floating ice; 26th, floating ice.

Wheeling, W. Va., 1st to 8th and 26th to 28th, floating ice; 1st to 4th, floating ice in Allegheny River.

Susquehanna River.-Lock Haven, Pa., 1st, river frozen; 10th, ice moved out; 16th, frozen.

-New York, N. Y., 15th, the first floating ice of the season Atlantic Coast .in the North and East rivers; 27th to 28th, floating ice in North and East rivers. Vineyard Haven, Mass., 25th, harbor frozen over, but ice broken by tide during day. Portland, Me., 23d to 28th, ice in harbor.

OBSERVATIONS ON THE GREAT LAKES.

Owing to the closing of navigation on the Great Lakes reports for the month of February from no vessels and from during the winter season the Weather Bureau has received only 10 U.S. Life Saving stations.

SUNSHINE AND CLOUDINESS.

received by the atmosphere above the cloud layer is very nearly constant from year to year, yet it varies with the day average cloudiness gives the observer's estimated percentage and the month. On the other hand, at the surface of the of clear sky, and these numbers are given in the last column earth, the distribution of sunshine and, therefore, the resultant of Table IV. On the average these personal estimated perheat with its meteorological and biological consequences depends mostly on the distribution of cloudiness.

Sunshine.—During the month an instrumental record of the amount of sunshine has been kept at 15 stations by means of the photographic sunshine recorder, and at 21 stations by means of the thermometric sunshine recorder. The both classes of instruments, yet this agreement needs an exresults of these observations are given in Table IV, which shows the actual sunshine received, on the average, for any hour of local mean time (not seventy-fifth meridian time) fact that the observer estimates the average area of the sky during the month; the actual sunshine is tabulated as a percentage of the greatest possible duration of sunshine.

between the hours of 11 a. m. and 1 p. m., are: Colorado from which is derived the percentage of duration of sunshine. Springs, Colo., 87.5; Denver, Colo., 87; Key West, Fla., 91; It is easily conceivable that a small bank of cloud slowly Portland, Me., 85.5; Tucson, Ariz., 91. The stations having moving across the sky might keep the instrument in its shadow the least percentage are: Cleveland, Ohio, 45.5; Galveston, Tex., 45.5; Memphis, Tenn., 42; New Orleans, La., 46; Portland, Oreg., 28.5; Savannah, Ga., 42.5.

The next to the last column of Table IV gives the general average sunshine for the whole month for all hours of daylight. The highest percentages are: Key West, Fla., 84; full sunshine. In this respect the various methods of record-Tucson, Ariz., 82; San Diego, Cal., 81. The lowest averages ing differ among themselves as though they had varying are: Galveston, Tex., 38; Memphis, Tenn., 39; New Orleans, degrees of sensitiveness: on clear and fair days the observer La., 35; Portland, Oreg., 23.

Although the quantity of sunshine, and probably of heat, based on numerous personal observations) are given for each Weather Bureau station in Table I. The complement of the centages are lower by 10.8 than the sunshine recorded by the thermometric registers and lower by 9.1 than the photographic registers.

Although the general average difference between the instrumental registers and personal records is about the same for planation. There is necessarily a systematic difference between the instrument and the observer, resulting from the covered by clouds at any moment, and his final percentage for the day, therefore, embraces the whole sky, whereas the The stations recording the largest percentage of sunshine instrument records essentially the visibility of the sun's disk, and cause a record of "no sunshine," whereas the observer viewing the whole sky would record that but a few tenths are covered by clouds. In general, in climates where but few clouds exist, an observer's record of one, two, or three tenths will almost invariably accompany an instrumental record of seems to be most sensitive to the presence of cloud, but the Cloudiness.—The number of clear and cloudy days and the instrument less so: in cloudy weather the observer seems average cloudiness between sunrise and sunset (the latter most sensitive to the presence of sunshine and the instru-